
An Analysis of Saline Water Intrusion into Coastal Nigeria

Dr. Amartya Kumar Bhattacharya

BCE (Hons.) (Jadavpur), MTech (Civil) (IIT Kharagpur), PhD (Civil) (IIT Kharagpur), Cert.MTERM (AIT Bangkok), CEng(I), FIE, FACCE(I), FISH, FIWRS, FIPHE, FIAH, FAE, MIGS, MIGS – Kolkata Chapter, MIGS – Chennai Chapter, MISTE, MAHI, MISCA, MIAHS, MISTAM, MNSFMFP, MIIBE, MICI, MIEES, MCITP, MISRS, MISRMTT, MAGGS, MCSI, MMBSI
Chairman and Managing Director, MultiSpectra Consultants,
23, Biplabi Ambika Chakraborty Sarani, Kolkata – 700029, West Bengal, INDIA.
E-mail:dramartyakumar@gmail.com

Abstract

The present paper gives an overview of saline water intrusion into coastal Nigeria. The paper places saline water intrusion into coastal Nigeria in the context of geology, hydrogeology, meteorology and land use of coastal Nigeria. The paper suggests ways and means to manage the problem of saline water intrusion into coastal Nigeria.

Introduction

Groundwater dynamics is the movement of groundwater through geological formations and it can be linked to different phenomena that occur in deep aquifers such as the upward advance of saline waters of geologic origin, surface waste discharges into shallow aquifers and the invasion of saline water into coastal aquifers.

In recent years, considerable interest has been evinced in the study of flow through porous media, because of their natural occurrence and importance in many engineering problems such as movement of water, oil and natural gas through the ground, saline water intrusion into the coastal aquifers, flow through packed towers in some chemical processes and filtration. Therefore, the physical understanding of flow through porous media is essential to scientists and engineers working in the related areas. The proposed work as reported herein is on movement of groundwater as related to saltwater intrusion into coastal aquifers and submarine discharge of freshwater into the sea. The mechanism responsible for this phenomenon involves the reduction or reversal of ground water gradients, which permits denser saline water to displace freshwater and vice-versa. This situation commonly occurs in coastal aquifer in hydraulic

continuity with the sea when pumping of wells disturbs the natural hydrodynamic balance.

Saline water intrusion in fresh groundwater takes place in the vicinity of coastal regions whenever saline water displaces or mixes with freshwater. This situation usually occurs in coastal regions having hydraulic continuity with sea when the pumping rate in the wells disturbs the natural hydrodynamic balance. Intrusion of saline water into freshwater coastal aquifer is likely to cause problem when such aquifer is tapped for domestic water supply or for irrigation.

The coast of Nigeria is located in West Africa, part of the Gulf of Guinea. The coastline of Nigeria stretches for about 850km, from Republic of Benin in the west to Cameroon in the east. It forms part of the West African groundwater region. River Niger discharges into the Gulf of Guinea via a delta as the major river, thereby, dividing the coastline into two parts. Other rivers discharge through estuaries either into lagoon or directly into the ocean. The surface strata of the coastline of Nigeria consists of unconsolidated coastal plain sands.

Lagoons, estuaries, creeks and delta dominate the Nigerian coastline. The vegetation of the area is mangrove swamp forest that is being cleared for wood and to give room for other economic activities. The western of the coastline is highly populated with total dependence on groundwater to meet domestic and industrial demands. Thousands of private boreholes have been drilled in the area to extract groundwater for drinking. Continuous pumping or heavy extractions of ground water have led to increment of chloride content in groundwater. The chloride/bicarbonate ratio is a very good indicator of saline water contamination in ground water. High chloride content in groundwater may also be caused by the presence of chloride containing minerals in the area or there are pollutants being discharged into groundwater. The major cause of saline water pollution into groundwater in this area is as a result of saline water intrusion. Upcoming occurs below wells as a result of continuous pumping of freshwater.

The entire coastline of Nigeria is divided into seven administrative units (Lagos, Ogun, Ondo, Delta, Bayelsa, Rivers and Akwa Ibom states).

Saline water intrusion is one of the most common forms of groundwater contamination in coastal areas (Bear, 1999). Saline water intrusion is the movement of salty water into an aquifer or surface water with the consequent mixing or displacement of fresh water. When groundwater is

pumped from aquifers that are in hydraulic connection with the sea, and the piezometric surface of the fresh water is lowered, the balance between the saline water / freshwater interface is disturbed, the hydraulic gradients so created induce sea water to encroach the freshwater aquifer (Domenico and Schwartz, 1998; Walton, 1970).

Saline water intrusion has been a major water resource problem in the urban coastal areas of the Niger Delta and it has therefore been and continues to be a focus of considerable research efforts over time. This is because a scientific understanding of the occurrence of salt and fresh waters in a particular coastal area is essential for the development and management of the water resource. Nigeria has a coastline that is about 1000km long with the Atlantic Ocean bordering eight States. These are: Rivers, Bayelsa, Akwa Ibom, Cross Rivers State, Delta, Ondo, Ogun and Lagos States. Potable water supply to inhabitants in some of the communities in the coastal belt (especially in the saline mangrove swamp) has been a major problem due to salt water intrusion. Water wells and boreholes drilled have been abandoned in many communities due to high salinity. They therefore depend on rain harvesting and purchasing water from water merchants. In Nigeria, there have been few studies aimed at assessing fresh water resources in coastal areas of the country. This forms the main thrust of the proposed research.

The quality of water is the main constraint of ground water in the coastal aquifers in Nigeria. Like all coastal aquifers worldwide, saline water intrusion into aquifers is the major source of quality impairment in the coastal aquifers in Nigeria. This is followed by pollution from spillages of crude and refined petroleum and lastly by leachates from municipal and other industrial wastes. Most of the confined aquifers in the coastal areas of Nigeria have high iron concentration such that treatment for the removal of iron has to be undertaken.

Saline water intrusion into unconfined and confined aquifers occurs in Niger Delta. In Coastal beach ridges or sandy islands within the saline mangrove belt, fresh water lens floating above saline water-bearing sands are found to occur in the unconfined aquifers (Oteri, 1990).

The growth of industrial development within the urban coastal areas of the Niger Delta and the attendant population explosion places a very heavy demand on the lean supply of fresh water. According to Akpokodje (1998) and Frank Briggs (2003)

“indiscriminate abstraction of ground water from the first aquifer has resulted in salt water intrusion in several coastal wells...”. This limits the supply of potable drinking water in the area, which can have detrimental effects on human health, wild life habitat and increase the cost of water treatment (Domenico and Schwartz, 1998).

Geography and Meteorology of Coastal Nigeria

In Nigeria, eight States are bounded to the south by the Atlantic Ocean. They include – from West to East – Ogun, Lagos, Ondo, Delta, Bayelsa, Rivers, Akwa Ibom and Cross River States. The States are contiguous and are in direct contact with the sea. They are located between latitude 7° and 4° 10' N and longitudes 2°30' and 8°30' East. There are a number of highly populated cities situated within the coastal area of these States which include-Port Harcourt, Warri, Yenagoa, Lagos, Uyo and Calabar.

The coastal States of Nigeria fall within humid tropical zone of the country. Rainfall is copious and lasts for eight to nine months of the year (mid March to early November) with mean annual value often exceeding 3000mm at the coastal fringes at Akasa and Brass. Seasonal changes of wet and dry are as a result of the interplay of two contrasting air masses – the moisture laden SW Monsoon Winds blowing into the country from the Atlantic Ocean and the dry North East Trade winds from the Sahara desert.

The two main rivers in the southwestern part of Nigeria are Ogun and Oshun which drain into the ocean.

Geology of Coastal Nigeria

Geologically, coastal Nigeria is made up of two sedimentary basins-the Dahomey Basin and the Niger Delta Basin separated by the Okitipupa Ridge.

The Dahomey Basin covers the southern areas of Lagos, Ogun, and Ondo States in Nigeria and stretches into the neighbouring countries of Benin, Togo, Ghana, and Ivory Coast.

The present study is centered on the Niger Delta area of Nigeria.

The geological formations of primary interest for the evaluation of the groundwater resources potential of the study area are the sedimentary formations deposited in the Niger Delta during the Eocene-Quaternary periods.

The stratigraphic sequence under consideration comprises, from oldest to youngest

Table 1: Stratigraphic Succession of the Niger Delta

SUB SURFACE			SURFACE OUTCROPS		
Youngest Known Age		Oldest Known Age	Youngest known Age		Oldest known Age
Recent	Benin Formation	Oligocene	Plio/Pliocene	Benin Formation	Miocene
Recent	Agbada Formation	Eocene	Miocene/Eocene	Ogwashi-Asaba Ameki Formation	Oligocene/Eocene
Recent	Akata Formation	Eocene	Lower Eocene Paleocene	Imo Shale Nsukka shale	Paleocene Maestrichtian
Equivalent not known			Maestrichtian Campanian Camp/Maest. Coniacian Santonian Turonian Albion	Ajali Fm Mamu Fm Nkporo Shale Agwu Shale Eze Aku Shale Asu River Group	Maestrichtian Campanian Santonian Turonian Albion

(Short and Stauble, 1967)

Table 2: Tertiary Correlation chart of the Niger Delta Succession and Outcrop Equivalent

AGE	SURFACE FORMATIONS	SUB SURFACE EQUIVALENTS	BROAD DEPOSITIONAL ENVIRONMENTS
PLIOCENE - RECENT	COASTAL PLAIN SANDS	BENIN FORMATION AFAM & QUA IBOE CLAY MEMBERS	CONTINENTAL
MIOCENE - RECENT	OGWASHI - ASABA FM IJEBU FM		
EOCENE - RECENT	AMEKI FORMATION ILARO & OSHOSHON FM	AGBADA FM	PARALIC
PALEOCENE - RECENT	IMO SHALE FM	AKATA FM	MARINE

formations, the Imo Shale Group, the Ameki Formation, the Benin Formation, and the Quaternary deposits as described by Amajor and Agbaire, (1989).

The Imo Shale Group is made up of marine clay, shale, and limestone. The formation is estimated to have a thickness of up to 1,000m and it outcrops in a belt more than 100km north of the study area. The formation, due to its impermeable texture, is believed to constitute the base of the groundwater aquifers under consideration in the study area.

The Ameki Formation overlies the Imo Shale Group. It comprises numerous alternations of marine shales and fine to coarse, very heterogeneous, coastal – deltaic sands and sandstones. The formation is estimated to have a thickness of up to 1,700m and is out-cropping in an east-west and north-west – south-east oriented belt, approximately 100km north of the study area. The groundwater resources potential of the formation is believed to be good (Tahal, 1998).

The present day Niger Delta is defined geologically by three subsurface sedimentary sequences consisting of Benin, Agbada and Akata Formations (Short and Stauble, 1967).

Benin Formation

The Benin Formation overlies the Ameki Formation. The formation is an extensive stratigraphic unit in the Southern Sedimentary Basin, with an average thickness of about 1,900m (Short and Stauble, 1967). The formation is recognised throughout the delta due to its few shale streaks, absence of brackish water and high percentage of sand.

The Benin Formation, which consists mainly of coastal plain sands, extends from the west across the whole Niger Delta area and southward beyond the present coastline. The sediments comprise yellow and white sands with pebbly gravels. The clays and sandy clays occur in lenses of 3m to as much as 10m, and they make the groundwater formation a multi- aquifer system. The formation is, a massive porous fresh water-bearing sandstone with localized thin beds. The formation is, due its coarse texture and huge outcrop area, believed to constitute a very good groundwater aquifer. The aquifer in this basin has a south-west gradient towards the Delta, and it thickens seawards in the same direction of groundwater movement. It is thus the most prolific aquifer in the region. It is overlain by the Quaternary deposits, which ranges in

thickness between 133 – 500ft (Etu – Efeotor and Akpokodje, 1990). The Quaternary deposit, which comprises recent deltaic sediments made up of sand, silt, and clay beds, overlies the Benin Formation in the swampy delta areas. The formations have a seaward dip resulting in confined aquifers.

The sandstones of the Benin Formation are coarse grained, locally fine – grained, poorly sorted and sub - angular to well rounded. The age of the Benin Formation lies between Oligocene and Recent. This formation constitutes the major aquifer in the Niger Delta area (Udom et al, 1999).

Whiteman (1982) notes that a formation such as the Benin Formation, deposited in a continental fluvial environment has a highly variable lithology, can be recognised because of its high sand percentage (70 – 100%), few minor shale intercalations and the absence of brackish water and marine faunas. To date, very little oil has been found in the Benin Formation (mainly minor oil shows).

Agbada Formation

The Agbada Formation underlies the Benin Formation and forms the second of the three strongly diachronous Niger Delta complex formations. It comprises mainly of alterations of sands, silts and shales in various thicknesses and proportions, indicative of cyclic sequences of off-lap cycles, better called off-lap rhythms (Weber, 1971).

The characteristic features of the sandstones here are poor sorting, calcareous matrix and shell fragment occurrences. The grain size varies from fine to coarse. The approximate total thickness of the Agbada Formation lies between 10,000ft and 12,000ft (Kogbe, 1976).

The top of the Agbada Formation is sandy whereas the bottom is shaley because it grades into the Akata shales gradually. The shales are denser at the base of the formation due to the compaction process. The sandy parts constitute the main hydrocarbon reservoirs in the delta oil fields. The shales constitute seals to the reservoirs and as such are very important because the formation is rich in hydrocarbon.

Paralic clastics of the Agbada Formation represent the true deltaic portions of the delta top-set and fluvio-deltaic environment (Short and Stauble, 1967; Weber and Dakoru, 1975). However, Whiteman (1982) notes that the Agbada Formation, as defined by Short and Stauble (1967), contains beds laid down in a variety of sub-environments

grouped together under the heading-paralic environment. The age of the Agbada Formation ranges from Eocene to Recent.

Akata Formation

This formation is the basal major time-transgressive lithological unit of the Niger Delta Complex. It is composed mainly of marine shales, but contains sands and silty beds which are thought to have been laid down as turbidites and continental slope channel fills. The shales are typically under-compacted and over-pressured, forming diapiric structures.

Whiteman (1982) contributes that the Akata Formation is rich in planktonic foraminifera, which indicate deposition on a shallow marine shelf environment.

The formation occurs between depths of 0-6000m below the Abgada Formation (Fig. 2) and ranges in age from Paleocene to Recent. It represents the pro – delta mega facies (Tahal, 1998).

Hydrogeology and Aquifers of the Study Area

In the study area, groundwater is abstracted from the Benin Formation, mainly from its upper section.

Lenses of silty clay of some few metres thickness have been recorded in the borehole penetrating the Benin Formation (Tahal 1998). To the south of Port Harcourt, belts rich in shales lying at a depth of 10m to 200m have been observed. These lenses create several sub-aquifers in the Benin Formation, the upper sub-unconfined, while the deeper aquifers range from leaky to confined and are isolated from the ground surface. The natural recharge comes mainly from the northern high coastal plain.

Generally, the sediments of the Benin Formation dip seawards at a low gradient. The upper section of these sediments is being utilised in the development of groundwater.

The sandy components, in most layers, are about 90% of the lithological sequences (Tahal, 1998). The size distribution of the sedimentary particulates does not vary significantly from place

Table 3: Lithological Variations of the Niger Delta

STRATIGRAPHICAL SEQUENCE			LITHOLOGY	MAXIMUM THICKNESS (m)
Quaternary	Alluvium in general		Gravel, sand, silt, clay	100
	Meander belt		Gravel, sand, clay	
	Wooded back-swamps and fresh water swamps		Silt, clay	
	Mangrove swamps		Fine sand, silt, clay	
	Lagoonal marshes		Silt, clay	
	Abandoned beach ridges		Sand, silt, clay	
	Sombreiro – Warri deltaic plain		Sand, clay, silt	
Tertiary	Miocene	Benin Formation (coastal plain sand)	Sand, coarse sand with clay and shales	2,000
		Ameki Formation	Clay, Sand, Shale	1,700
	Eocene	Imo Shale Group	Clay, sand, shale, limestone	1,000

(Kampsax – Krüger, 1985)

to place in the study area. Table 3 shows the lithological variations of the Niger Delta. There is a steady gradation in quantity between coarse sands and clay beds. Locally to some extent, the clay beds separate hydraulically between the sub-aquifers.

Groundwater abstracted from the upper Benin Formation is characterised by low p^H (acidity), high carbondioxide (CO_2) content, and hence it is corrosive and soft. In addition, the groundwater contains high concentrations of iron and therefore requires treatment.

The water tables of all sub-aquifers penetrated by boreholes are relatively shallow and range between 5m to 15m where hydrostatic water levels increase with the depth (Tahal, 1998).

According to Etu – Efeotor and Odigi (1983), three main zones have been differentiated as follows:

- (I) a northern zone consisting shallow aquifer of predominantly continental deposits.
- (ii) a transitional zone of marine and continental materials.
- (iii) a coastal zone of predominantly marine deposits.

Aquifer distributions in the Niger Delta are controlled by the geology. Table 4 is the summary of the properties and behaviour of hydrostratigraphic units in the Niger Delta Basin. In most parts of the Niger Delta, including the study area, a multi – aquifer system is encountered and the aquifer lies within the arenaceous Benin Formation.

The depth to water table of the Benin Formation ranges between 3m – 5m below ground level (Offodile, 1982).

Ngah (1990) also identified three main aquifer zones in the Niger Delta Viz:

- (i) an upper unconfined aquifer extending throughout the Benin Formation with its thickness ranging between 15m – 80m while the Static Water Level (SWL) varies between 4m and 21 m.
- (ii) a middle aquifer system, semi – confined and consisting of thick medium to coarse grained, sometimes pebbly sands with thin clay lenses. Its thickness varies between 30m – 60m

Table 4: Summary of Properties and Behaviour of Hydrostratigraphic Units in the Niger Delta Basin

Hydrogeol- ogical basin	Hydrostratig- raphic units	Lithologic details	Aquifer type and characteristics	Water quality	Economic importance
Niger Delta Basin	Alluvial plains aquifer	Sands, clays, silt	Unconfined	Saline water	Domestic MunicipalInd ustrial uses
	Meander Beltaquifer	Sands, gravel, clays		Corrosion	
	Mangrove swamp aquifer	Sands, clay, swamps	Good, water table is confined	Iron – rich saline water intrusion	
	Abandoned Ridges aquifer, Sombreiro	Sands and pebbles, yellow sands, clays	Poor aquifer		
	Benin aquifer	Sand, clays	Prolific		
	DeltaOgwashi – AsabaAmeki aquifer	Clays, shale, lignite, silty sand, clay siltstone	Aquitard	Problematic with iron and salt water encroachment	
	Imo shalesDeep oceanicAgbada – AkpataShales	Shalestone, claystone lenses of sands	Aquitard Aquitard	Low Ph	

- (iii) a lower aquifer system that extends from 220m – 300m and consists of coarse – grained sands and gravels with some interlayer clay.

The majority of the groundwater wells abstract water from the first and second aquifers (<100m deep). The very few industrial and municipal ground water supply wells tap deeper aquifers.

Aquifer Recharge in the Study Area

Precipitation is the main source of recharge to aquifers. The rate of recharge depends to a large extent on the infiltration capacity of the soil, evaporation rate and the overland drainage characteristics. In places where sandy clay forms a part of soil layers, recharge usually occurs mainly through a distant outcrop of the porous formation and partially through the lateritic sand.

The northern and the southern movement of air masses and pressure belts characterise the rainfall regime of the coastal plain. The amount of rainfall decreases inland from the coast. The isohyets run parallel to the coast up to a line through Ahoada-Degema-Port Harcourt. From this line inland, the decrease of rainfall is much slower.

According to the lithologic profile of some boreholes, in some places there are few metres of lateritic sandy clay under the soil. Tahal (1998) observed that about 30% to 40% of the yearly average of rainfall (2,280mm) could infiltrate and recharge the Benin Formation. To the north, sandy and porous outcrops exist, where the replenishment can be 60% to 70% of rainfall.

The estimated annual recharge to the aquifer in the study area from the Northern High Coastal Plain is about 100mcm to 150mcm (Tahal, 1998).

Problems Affecting the Coastal Aquifer Ecosystem in Nigeria

The major constraint to the water resource in the aquifers of the coastal Nigeria is quality. The saline water intrusion is both natural and man-made. For the natural saline water intrusion, the problem is exacerbated by:

- The lack of willingness to carry out necessary studies or utilise such studies when carried out
- Lack of appreciation of the need for water resources assessment and management of ground water resources of the coastal aquifers resulting in over-exploitation of the aquifers.

The activities of man have also led to increased saline water intrusion through:

- Uncontrolled development of both unconfined and confined aquifers especially in Port Harcourt Metropolis, Lagos, Warri and Bonny. In Lagos which is the commercial capital of Nigeria, the problem is particularly acute as many boreholes which were producing fresh water after drilling become salty a few months after especially in Ikoyi, Victoria Island and Apapa.
- Lack of proper sealing of disused boreholes or those abandoned due to salt water intrusion.
- The groundwater in most of the confined aquifers in coastal Nigeria is corrosive and casing corrosion is a major source of borehole failures. There is a need to determine the best material for casing and screens and also best completion techniques that will prevent saline water ingress into fresh water yielding boreholes especially in situations where the fresh water aquifers underlie saline water bearing sands.
- Sea water level rise (as a result of global warming); building of dams in upstream areas of the rivers leading to low flows and decrease in river sediments to the coast by the rivers have all led to increased coastal erosion.
- Construction of drainage canals, transportation canals
- Urbanisation – In the Lagos area for example. The rate of Urbanisation is so high that recharge area have been converted to concrete zones, while swamps and streams are being reclaimed and turned into cities. This decreases the amount of recharge to the aquifers, thereby increasing saline water intrusion.
- Oil spillage from both upstream and downstream operations of the oil industry is the next major problem affecting both the aquifers and the environment of coastal areas in general.
- Finally, disposal of wastes from industries and municipal areas is a threat to coastal aquifer ecosystem.

Management of the Problem

The control of saline water intrusion demands knowledge of the hydraulic conditions within aquifer; it also demands knowledge of the source of

the saline water. It is therefore necessary to identify the extent of the problem and to assess the behaviour of the saline water body under various conditions of recharge and discharge, such that efficient water resources management plan can be implemented. The optimum solution to the problem of saline water intrusion is prevention, by which the encroachment of saline water is controlled to an acceptable degree. But in many cases, the problem is a legacy of the past, therefore, management must concentrate on minimizing further intrusion, and/or reducing the extent of the existing saline water, it may be that the aquifer in question is too badly polluted and so reclamation may be the only viable option. In extreme cases if the resource for potable water supply, it may be abandoned, although the water may still be utilised for certain industrial or agricultural applications.

References

- Amajor, L.C and Agbaire, D.W (1989). Depositional History of the Reservoir Sandstones, Akpor and Apará Oilfields, Eastern Niger Delta, *Nigeria Journal of Petroleum Geology*. Vol. 12, No.4 pp. 453-464.
- Bear, J; Cheng, A.H.D., Sorek, S., Ouazar, D. & Herrer, T. (1999). J. Bear (ed) Saltwater intrusion in Coastal Aquifers – Concepts, Methods and Practices, in Theory and Application in porous Media, Kluwer Academic Publishers. Dordrecht, P 625.
- Domenico, P.A and Schwartz, F.W (1988). *Physical and Chemical Hydrogeology*, John Wiley and Sons Inc., New York.
- Etu – Efeotor, J.O and Akpokodje, G.E (1990). Aquifer Systems of the Niger Delta. *Journal of Mining and Geology*. Vol. 26, No.2, pp264 – 266.
- Etu – Efeotor, J.O and Odigi, M.I (1981). Water Supply Problems in the Eastern Niger Delta. *Journal of Mining and Geol.*, Vol. 20, No.1 pp 182 - 192.
- Frank-Briggs, I.N (2003). *The Geology of some Island Towns in the Eastern Niger Delta, Nigeria*. An Unpublished Ph.D Thesis Submitted to the Department of Geology, University of Port Harcourt.
- Kogbe, C.A (1976). *The Cretaceous and Paleocene Sediments of Southern Nigeria*; Elizabethan Publishing Coy; Lagos. Pp 237 – 252
- Kampsax – Krüger (1985). *Final Report on Hydrological Site Studies for Water Well Development, Phase 1*, p.118.
- Ngah, S.A (1990). *Groundwater Resource Development in the Niger Delta: Problems and prospects*. 6th IAEC Congress.
- Offodile, M.E (1982). The Problems of Water Resources Management in Nigeria; *Journal of Mining and Geology, 20th Anniversary Edition*, Vol. 19, No. 1.
- Oteri, A.U (1990). Delineation of sea water intrusion in a coastal beach ridge of Forcados. *Journal of Mining and Geology* 26 (2) 225 – 229.
- Short, K.C and Stauble, A.J (1967). Outline of the Geology of the Niger Delta. *AAPG Bull.* Vol 51., pp 761 – 779.
- Tahal Consultant Engineers Ltd (1998). *Final Report on Multi – State Water Supply Project, Feasibility Study of Rivers State Capital (Port Harcourt and Environs and Selected Urban Communities)* pp. 2-4.
- Udom, G.J, Etu-Efeotor, J.O and Esu. E.O (1999). Hydrochemical Evaluation of Groundwater in parts of Port Harcourt and Tai-Elime Local Government Area, Rivers State: *Global Journal of Pure and Applied Sciences*, Vol.1.5, pp. 546 – 552.
- Weber, K.J (1971). Sedimentological Aspects of Oil Fields in the Niger Delta. *Journal of Environmental Geology*, Minbouw, Vol. 50, No.3, pp 559-576
- Weber, K.J and Dakoru, E (1975). *Petroleum Geology of the Niger Delta*. Ninth World Petroleum Congress. pp. 109-221.
- Whiteman, A (1982). *Nigeria: Its Petroleum Resources and Potentials*. Graham and Trotman. Vol. 1, pp 63-78.
- Walton, W.C., (1970). *Groundwater Resource Evaluation*, McGraw – Hill. P. 375.

● POWER & CONTROLS ● PUMPING STATION & PIPELINE ● SECURITY & FIRE FIGHTING SYSTEM
● INDUSTRIAL CONSTRUCTION & CIVIL ● INSTRUMENTATION ● MAINTENANCE/UTILITY SERVICE

Please Contact:

**LOGIC
NODES**

23E, Fern Road, Kolkata-700 019 ● Phone : 033-2460-3689, 2460-3155 ● Telefax : 033-2440-2739
E-mail: ulogdeta1@gmail.com, ulogdeta@yahoo.com ● Website : www.logicnodes.in