

CORAL REEFS

Plastic waste associated with disease on coral reefs

Joleah B. Lamb,^{1,2,3*} Bette L. Willis,^{2,3} Evan A. Fiorenza,^{1,4} Courtney S. Couch,^{1,5,6} Robert Howard,⁷ Douglas N. Rader,⁸ James D. True,⁹ Lisa A. Kelly,^{3,10} Awaludinnoer Ahmad,^{11,12} Jamaluddin Jompa,¹² C. Drew Harvell¹

Plastic waste can promote microbial colonization by pathogens implicated in outbreaks of disease in the ocean. We assessed the influence of plastic waste on disease risk in 124,000 reef-building corals from 159 reefs in the Asia-Pacific region. The likelihood of disease increases from 4% to 89% when corals are in contact with plastic. Structurally complex corals are eight times more likely to be affected by plastic, suggesting that microhabitats for reef-associated organisms and valuable fisheries will be disproportionately affected. Plastic levels on coral reefs correspond to estimates of terrestrial mismanaged plastic waste entering the ocean. We estimate that 11.1 billion plastic items are entangled on coral reefs across the Asia-Pacific and project this number to increase 40% by 2025. Plastic waste management is critical for reducing diseases that threaten ecosystem health and human livelihoods.

Outbreaks of disease on coral reefs threaten one of the most biodiverse ecosystems on the planet (1), jeopardizing the U.S. \$375 billion in goods and services that they provide to people each year through fisheries, tourism, and coastal protection (2). Plastic waste can host pathogens that are frequently implicated as triggers of disease outbreaks on coral reefs (3–9). For example, microbial communities colonizing polypropylene marine debris were dominated by the genus *Vibrio* (10), an opportunistic pathogenic bacteria of a globally

devastating group of coral diseases known as white syndromes (11). Although an estimated 4.8 million to 12.7 million metric tons of plastic waste enter the ocean in a single year (12), the resulting influence on disease susceptibility in the marine environment is unknown. Microbial rafting on plastic debris has been shown to strongly control surface longevity (13) and is highest in tropical regions near the equator compared with more polar regions (14), suggesting that coral reef ecosystems could have high levels of colonized plastic waste.

We surveyed 159 coral reefs spanning eight latitudinal regions from four countries in the Asia-Pacific for plastic waste and evaluated the influence of plastic on diseases that affect keystone reef-building corals (15) (benthic area = 12,840 m²) (Fig. 1). The Asia-Pacific region contains 55.5% of global coral reefs (2) and encompasses 73.0% of the global human population residing within 50 km of a coast (12) (table S1). Overall, we documented benthic plastic waste (defined as an item with a diameter >50 mm) on one-third of the coral reefs surveyed, amounting to 2.0 to 10.9 plastic items per 100 m² of reef area [95% confidence interval (CI), $n = 8$ survey regions]. The number of plastic items observed on each reef varied markedly among countries, from

¹Department of Ecology and Evolutionary Biology, Cornell University, Ithaca, NY 14853, USA. ²Australian Research Council Centre of Excellence for Coral Reef Studies, James Cook University, Townsville, Queensland 4811, Australia. ³College of Science and Engineering, James Cook University, Townsville, Queensland, Australia. ⁴School of Aquatic and Fishery Sciences, University of Washington, Seattle, WA 98195, USA. ⁵Hawaii Institute of Marine Biology (HIMB), University of Hawaii at Manoa, Kaneohe, HI 96744, USA. ⁶Ecosystem Sciences Division, National Oceanic and Atmospheric Administration (NOAA) Pacific Islands Fisheries Science Center, Honolulu, HI 96818, USA. ⁷Marine Programme, Fauna & Flora International, Yangon, Myanmar. ⁸Oceans Program, Environmental Defense Fund, New York, NY 10010, USA. ⁹Center for Biodiversity in Peninsular Thailand, Prince of Songkla University, Hat Yai, Songkhla, Thailand. ¹⁰Fish Ecology and Conservation Physiology Laboratory, Department of Biology and Institute of Environmental Science, Carleton University, Ottawa, Ontario, Canada. ¹¹The Nature Conservancy, Raja Ampat Field Office, North Sorong, West Papua, Indonesia. ¹²Faculty of Marine Science and Fisheries, Hasanuddin University, Makassar, South Sulawesi, Indonesia. *Corresponding author. Email: joleah.lamb@cornell.edu




Fig. 1. Estimated plastic debris levels on coral reefs. (A) Modeled association between plastic debris on coral reefs from surveys of 159 reefs in eight regions [red squares in (B)] from 2011–2014 and estimated levels of mismanaged plastic waste (thousand metric tons), assuming that 25% of waste entered the ocean in 2010 from human populations living within 50 km of the coast for each country (12). Reef locations can be found in table S14. The Asia-Pacific region encompasses 9 of 10 countries with the highest global levels of estimated mismanaged plastic waste entering the ocean (table S3). Gray shading represents the upper and lower 95%

CI for the model. (B and C) Modeled plastic debris levels on coral reefs (100 m²), as projected using the association between estimated mismanaged plastic waste entering the ocean in 2010 for each sovereign country (12) and plastic debris surveys between 2011–2014 [shown in (A)]. The color scale represents the minima and maxima model estimates of mismanaged plastic waste on coral reefs from 2010 (table S5). Projections of plastic debris on coral reefs for Indonesia and China in 2025 were set to the maxima from 2010, owing to the limitations of the model range. Countries without a coastline are shown in white.

maxima in Indonesia [25.6 items per 100 ± 12.2 m² (here and elsewhere, the number after the \pm symbol denotes SEM)] to minima in Australia (0.4 items per 100 ± 0.3 m²) (table S2).

Terrestrially derived pollutants have been implicated in several disease outbreaks in the ocean (16). However, no studies have examined the influence of plastic waste on disease risk in a marine organism. In this work, we visually examined 124,884 reef-building corals for signs of tissue loss characteristic of active disease lesions (15) (fig. S1). We found plastic debris on 17 genera from eight families of reef-forming corals. When corals were not in contact with plastic debris, the likelihood of disease was $4.4 \pm 0.2\%$ across all eight regions (range = 2.8 to 8.4%, generalized linear mixed model, likelihood ratio test among regions: $\chi^2_7 = 10.382$, $P = 0.168$) (Fig. 2A). In contrast, in the presence of plastic debris, the likelihood of disease occurrence in corals increased significantly by more than a factor of 20 to $89.1 \pm 3.2\%$ (generalized linear mixed model: z score = 27.24, $P < 0.001$, $n = 331$ transects) (Fig. 2B and table S3).

Human population size in coastal regions and the quality of waste management systems largely determine which countries contribute the greatest plastic loads entering the ocean, given that an estimated 80% of marine plastic debris originates from land (12). Accordingly, we modeled the relationship between our documented levels of plastic debris on coral reefs ($n = 437$ transects from Australia, Myanmar, Thailand, and Indonesia) and Jambek *et al.*'s estimated levels of mismanaged plastic waste entering the ocean (12) from these four countries in 2010 (15) (generalized linear mixed model: Akaike information criterion = 662.3, $z = 3.95$, $P < 0.001$) (Fig. 1A, fig. S2, and table S4). Our model encompasses the range of mismanaged plastic waste entering the ocean introduced by coastal populations from 15 of the 17 (88%) sovereign countries in the Asia-Pacific region (maximum = 804,214 metric tons, minimum = 3472 metric tons), of which 9 are among the top 10 plastic-polluting countries globally (table S5). Assuming that improvements in waste management infrastructure did not occur during our survey period (2011–2014) and that the plastic waste emanated from adjacent terrestrial point sources, we estimate that levels of plastic debris on coral reefs for each country in the Asia-Pacific ranged from 0.9 to 26.6 plastic items per 100 m² in 2010 (95% CI) (Fig. 1B and table S5). This amounts to an estimated 11.1 billion items of plastic on coral reefs across the Asia-Pacific (95% CI = 1.2 billion to 105.5 billion items, $n = 15$ countries), which is likely underestimated owing to the exclusion of China and Singapore because they fall outside of the model range (table S5).

By 2025, the cumulative quantity of plastic waste potentially entering the marine environment from land is predicted to increase by one order of magnitude (12). Using this projection and assuming that the area encompassed by coral reefs remains constant, we estimate that 15.7 billion plastic items will be entangled on coral reefs across the Asia-Pacific by 2025 (the “business-as-usual” sce-




Fig. 2. Plastic waste influences disease susceptibility of reef-building corals. (A and B) Box (median and 50% quantile) and whisker (95% quantile) plots of coral disease likelihood for each of eight regions in four countries in the Asia-Pacific when no plastic waste is present (A) ($n = 362$ transects) and when plastic waste is present (B) ($n = 75$ transects). The red line represents the mean and the light red bar denotes ± 1 SEM across all eight regions. Boxes are shaded according to model estimates of plastic debris on coral reefs per 100 m² from Fig. 1C. MYAN, Myanmar; THAIL, Thailand; INDO, Indonesia; AUST, Australia.




Fig. 3. Reef-building corals with plastic debris have different disease assemblages than corals without plastic debris. Multivariate spatial representation of the relative abundance and composition of coral disease assemblages, as determined by a principal coordinates analysis (PCoA) ($n = 75$ paired transects). Vectors for each group illustrate the median spatial distance within the group. Disease assemblages represent six diseases—skeletal eroding band, white syndromes, black band, growth anomalies, brown band, and atramentous necrosis—recorded commonly across the globe.

nario for global infrastructure: 95% CI = 1.7 billion to 149.2 billion items) (Fig. 1C and table S5). According to our model, the predicted geographic distribution of plastic debris on coral reefs does not change substantially between now and 2025 (Fig. 1, B and C), but the disparity in quantities of accumulated plastic waste between developing and industrialized countries grows considerably. For example, plastic debris on coral reefs increases by only ~1% in high-income countries such as Australia but nearly doubles in a similarly populated low-income country such as Myanmar (Fig. 1C and table S5).

Comparative analyses of disease prevalence among different diseases in the presence versus absence of plastics can offer insights into potential mechanisms that increase disease susceptibility in corals. Reef-building corals in contact with plastic debris were affected by four of six common diseases globally (17), whereas corals

without plastic debris were affected by all six diseases but at much lower prevalence levels (table S6). Disease assemblages on reef-building corals differ distinctly when contact with plastic waste is present versus absent, as visualized by a principal coordinates ordination analysis (15) (permutational multivariate analysis of variance: $F = 11.86$, $P < 0.001$, $n = 75$ paired transects) (Fig. 3). In particular, three key diseases associated with rapid coral mortality increased markedly when plastic debris was in contact with coral tissues: Skeletal eroding band disease increased from $1.2 \pm 0.1\%$ to $43.9 \pm 5.1\%$ (increased likelihood = 24%), white syndromes increased from $1.9 \pm 0.2\%$ to $19.0 \pm 4.0\%$ (increased likelihood = 17%), and black band disease increased from $0.6 \pm 0.1\%$ to $14.7 \pm 3.9\%$ (increased likelihood = 5%) (tables S7 to S10).

Given the widespread distribution of plastic debris on coral reefs and the consequent increased




Fig. 4. Coral morphological complexity influences risk to plastic debris and disease. (A to D) Posterior probability density functions of coral species grouped into three broad morphological classifications. Structural complexity is determined by coral species (18); see table S11 for classifications. Minimum,

maximum, and peak values are shown for each structural complexity classification group: massive (dark red), branching (medium red), and tabular (light red). For ease of comparison, the inset in (C) represents the likelihood of disease without plastic debris [as shown in (B)].

likelihood of coral mortality from disease, we evaluated the potential for plastic debris and disease to affect structural complexity provided by habitat-forming corals. The structural complexity formed by corals underpins the availability of microhabitats for coral reef-associated organisms (18). We grouped coral species into three broad classifications based on the increasing structural complexity of their colony morphologies (massive < branching < tabular) (table S11) and determined that plastic debris is eight times more likely to affect reef corals with greater structural complexity (tabular and branching versus massive, $n = 348$ transects; posterior probability functions) (Fig. 4A and table S12). Massive coral morphologies are less likely to maintain contact with plastic debris; however, they exhibit the greatest increase in disease risk when this occurs (likelihood is increased by 98%) (Fig. 4, B to D, and table S13).

Our study shows that plastic debris increases the susceptibility of reef-building corals to disease. Plastics are a previously unreported correlate of disease in the marine environment. Although the mechanisms remain to be investigated, the influence of plastic debris on disease development may differ among the three main global diseases that we observed. For example, plastic debris can cause physical injury and abrasion to coral tissues by facilitating invasion of pathogens (19) or by exhausting resources for immune system function during wound-healing processes (20). Experimental studies show that artificially inflicted wounds to corals are followed by the establishment of the ciliated protozoan *Halofolliculina corallasia*, the causative agent of skeletal eroding band disease (21). Plastic debris could also directly introduce resident and foreign pathogens or may indirectly alter beneficial microbial symbionts. Cross-ocean bacterial colonization of polyvinylchloride (PVC) is dominated by *Rhodobacteriales* (22), a group of potentially opportunistic pathogens associated with outbreaks of several coral diseases (23). Additionally, recent studies have

shown that experimental shading and low-light microenvironments can lead to anoxic conditions favoring the formation of polymicrobial mats characteristic of black band disease (24).

By disproportionately reducing the composition or abundance of structurally complex reef-building coral species through disease, widespread distribution of plastic waste may have negative consequences for biodiversity and people (25). For example, on coral reefs, the loss of structural habitat availability for reef organisms has been shown to reduce fishery productivity by a factor of 3 (26).

Climate-related disease outbreaks have already affected coral reefs globally and are projected to increase in frequency and severity as ocean temperatures rise (27). With more than 275 million people relying on coral reefs for food, coastal protection, tourism income, and cultural importance (2), moderating disease outbreak risks in the ocean will be vital for improving both human and ecosystem health. Our study indicates that decreasing the levels of plastic debris entering the ocean by improving waste management infrastructure is critical for reducing the amount of debris on coral reefs and the associated risk of disease and structural damage.

REFERENCES AND NOTES

1. C. D. Harvell *et al.*, *Science* **296**, 2158–2162 (2002).
2. L. Burke, K. Reytar, M. Spalding, A. Perry, *Reefs at Risk Revisited* (World Resources Institute, 2011).
3. I. V. Kirstein *et al.*, *Mar. Environ. Res.* **120**, 1–8 (2016).
4. S. Oberbeckmann, M. G. Löder, M. Labrenz, *Environ. Chem.* **12**, 551–562 (2015).
5. C. A. De Tender *et al.*, *Environ. Sci. Technol.* **49**, 9629–9638 (2015).
6. K. M. Morrow, R. Ritson-Williams, C. Ross, M. R. Liles, V. J. Paul, *PLOS ONE* **7**, e44859 (2012).
7. M. M. Nugues, G. W. Smith, R. J. Hoodonk, M. I. Seabra, R. P. Bak, *Ecol. Lett.* **7**, 919–923 (2004).
8. J. E. Smith *et al.*, *Ecol. Lett.* **9**, 835–845 (2006).
9. M. J. Sweet, J. C. Bythell, M. M. Nugues, *PLOS ONE* **8**, e69717 (2013).
10. E. R. Zettler, T. J. Mincer, L. A. Amaral-Zettler, *Environ. Sci. Technol.* **47**, 7137–7146 (2013).
11. D. Bourne, T. Ainsworth, F. Pollock, B. Willis, *Coral Reefs* **34**, 233–242 (2015).

12. J. R. Jambeck *et al.*, *Science* **347**, 768–771 (2015).
13. F. M. Fazey, P. G. Ryan, *Environ. Pollut.* **210**, 354–360 (2016).
14. D. K. Barnes, *Nature* **416**, 808–809 (2002).
15. Materials and methods are available as supplementary materials.
16. S. Altizer, R. S. Ostfeld, P. T. J. Johnson, S. Kutz, C. D. Harvell, *Science* **341**, 514–519 (2013).
17. J. B. Lamb, D. H. Williamson, G. R. Russ, B. L. Willis, *Ecology* **96**, 2555–2567 (2015).
18. L. E. Richardson, N. A. J. Graham, A. S. Hoey, *Sci. Rep.* **7**, 7557 (2017).
19. J. B. Lamb *et al.*, *Philos. Trans. R. Soc. London Ser. B* **371**, 20150210 (2016).
20. L. D. Mydlarz, L. E. Jones, C. D. Harvell, *Annu. Rev. Ecol. Evol. Syst.* **37**, 251–288 (2006).
21. C. A. Page, B. L. Willis, *Coral Reefs* **27**, 257–272 (2008).
22. H. Dang, T. Li, M. Chen, G. Huang, *Appl. Environ. Microbiol.* **74**, 52–60 (2008).
23. N. Soffer, J. Zaneveld, R. Vega Thurber, *Environ. Microbiol.* **17**, 1203–1218 (2015).
24. M. S. Glas, Y. Sato, K. E. Ulstrup, D. G. Bourne, *ISME J.* **6**, 1526–1534 (2012).
25. B. Worm *et al.*, *Science* **314**, 787–790 (2006).
26. A. Rogers, J. L. Blanchard, P. J. Mumby, *Curr. Biol.* **24**, 1000–1005 (2014).
27. J. Maynard *et al.*, *Nat. Clim. Chang.* **5**, 688–694 (2015).

ACKNOWLEDGMENTS

We thank S. Atto, S. Beveridge, F. J. Pollock, G. Torda, A. Tracy, A. Wenger, C. Wood, S. Yusuf, and S. Piromvaragorn for their assistance. This research was supported by The Nature Conservancy NatureNet Science Fellowship, an AIMS@JCU (Australian Institute of Marine Science at James Cook University) Postgraduate Scholarship, the NSF Ecology of Infectious Marine Disease Research Coordination Network [Division of Ocean Sciences (OCE) award 1215977], the Capturing Coral Reef and Ecosystem Related Services (CCRES) Project funded by the Global Environment Facility and the World Bank (project ID P123933), NOAA/National Ocean Service (NOS)—HIMB agreement code MOA-2009-039, Australian Research Council (grant CE0561435), and the Environmental Defense Fund Innovation for Impact partnership with the Atkinson Center for a Sustainable Future at Cornell University. Surveys in Australia were conducted under Great Barrier Reef Marine Park Authority permits G10/33393.1 and G12/35232.1. All data and code to understand and assess the conclusions of this research are available in the main text, supplementary materials, and via the Dryad Digital Repository (<https://doi.org/10.5061/dryad.mp480>).

SUPPLEMENTARY MATERIALS

www.sciencemag.org/content/359/6374/460/suppl/DC1
Materials and Methods
Figs. S1 and S2
Tables S1 to S14
References (28–30)

31 October 2017; accepted 21 December 2017
10.1126/science.aar3320

Plastic waste associated with disease on coral reefs

Joleah B. Lamb, Bette L. Willis, Evan A. Fiorenza, Courtney S. Couch, Robert Howard, Douglas N. Rader, James D. True, Lisa A. Kelly, Awaluddinnoer Ahmad, Jamaluddin Jompa and C. Drew Harvell

Science **359** (6374), 460-462.
DOI: 10.1126/science.aar3320

Corals wrapped in plastic

Coral reefs provide vital fisheries and coastal defense, and they urgently need protection from the damaging effects of plastic waste. Lamb *et al.* surveyed 159 coral reefs in the Asia-Pacific region. Billions of plastic items were entangled in the reefs. The more spikey the coral species, the more likely they were to snag plastic. Disease likelihood increased 20-fold once a coral was draped in plastic. Plastic debris stresses coral through light deprivation, toxin release, and anoxia, giving pathogens a foothold for invasion.

Science, this issue p. 460

ARTICLE TOOLS

<http://science.sciencemag.org/content/359/6374/460>

SUPPLEMENTARY MATERIALS

<http://science.sciencemag.org/content/suppl/2018/01/24/359.6374.460.DC1>

REFERENCES

This article cites 27 articles, 6 of which you can access for free
<http://science.sciencemag.org/content/359/6374/460#BIBL>

PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)